

## Think | BIG

### Behavior Profile References: Test

1. Almond D, Madanitsa M, Mwapasa V, et al. Provider and user acceptability of intermittent screening and treatment for the control of malaria in pregnancy in Malawi. *Malar J.* 2016;15:574. <https://malariajournal.biomedcentral.com/articles/10.1186/s12936-016-1627-5>.
2. Altaras R, Nuwa A, Agaba B, Streat E, Tibenderana JK, Strachan CE. Why do health workers give anti-malarials to patients with negative rapid test results? A qualitative study at rural health facilities in western Uganda. *Malar J.* 2016;15:23. <https://malariajournal.biomedcentral.com/articles/10.1186/s12936-015-1020-9>.
3. Amoakoh-Coleman M, Agyepong IA, Kayode GA, Grobbee DE, Klipstein-Grobusch K, Ansah EK. Public health facility resource availability and provider adherence to first antenatal guidelines in a low resource setting in Accra, Ghana. *BMC Health Serv Res.* 2016;16:505. <https://bmchealthservres.biomedcentral.com/articles/10.1186/s12913-016-1747-1>.
4. Banchani E, Tenkorang EY. Implementation challenges of maternal health care in Ghana: the case of health care providers in the Tamale Metropolis. *BMC Health Serv Res.* 2014;14:7. <https://bmchealthservres.biomedcentral.com/articles/10.1186/1472-6963-14-7>.
5. Boadu NY, Amuasi J, Ansong D, Einsiedel E, Menon D, Yanow SK. Challenges with implementing malaria rapid diagnostic tests at primary care facilities in a Ghanaian district: a qualitative study. *Malar J.* 2016;15:126. <https://www.ncbi.nlm.nih.gov/pubmed/26921263>.
6. Cundill B, Mbakilwa H, Chandler CI, et al. Prescriber and patient-oriented behavioural interventions to improve use of malaria rapid diagnostic tests in Tanzania: facility-based cluster randomised trial. *BMC Med.* 2015;13. <https://bmcmmedicine.biomedcentral.com/articles/10.1186/s12916-015-0346-z>.
7. Chandler CIR, Meta J, Ponzio C, et al. The development of effective behaviour change interventions to support the use of malaria rapid diagnostic tests by Tanzanian clinicians. *Implement Sci.* 2014;9:83. <https://www.ncbi.nlm.nih.gov/pubmed/24969367>.
8. Diggle E, Asgary R, Gore-Langton G, et al. Perceptions of malaria and acceptance of rapid diagnostic tests and related treatment practises among community members and health care providers in Greater Garissa, North Eastern Province, Kenya. *Malar J.* 2014;13:502. <https://www.ncbi.nlm.nih.gov/pubmed/25519710>.
9. Febir LG, Baiden FE, Agula J, et al. Implementation of the integrated management of childhood illness with parasitological diagnosis of malaria in rural Ghana: health worker perceptions. *Malar J.* 2015;14:174. <https://malariajournal.biomedcentral.com/articles/10.1186/s12936-015-0699-y>.
10. Hill E, Hess R, Aborigo R, et al. "I don't know anything about their culture": the disconnect between allopathic and traditional maternity care providers in rural northern Ghana. *Afr J Reprod Health.* 2014;18(2):36-45. <https://www.ncbi.nlm.nih.gov/pubmed/25022140>.
11. Johansson EW. Beyond 'test and treat' – malaria diagnosis for improved pediatric fever management in sub-Saharan Africa. *Glob Health Action.* 2016;9(1):31744. <https://www.tandfonline.com/mutex.gmu.edu/doi/abs/10.3402/gha.v9.31744>.
12. Kwarteng A, Asante KP, Abokyi L, et al. Provider compliance to artemisinin-based combination therapy at primary health care facilities in the middle belt of Ghana. *Malar J.* 2015;14:361. <https://malariajournal.biomedcentral.com/articles/10.1186/s12936-015-0902-1>.

13. Nwameme AU, Phillips JF, Adongo PB. Compliance with Emergency Obstetric Care Referrals Among Pregnant Women in an Urban Informal Settlement of Accra, Ghana. *Matern Child Health J.* 2014;18(6):1403-1412. <https://www.ncbi.nlm.nih.gov/pubmed/24178157>.
14. Oduro-Mensah E, Kwamie A, Antwi E, et al. Care Decision Making of Frontline Providers of Maternal and Newborn Health Services in the Greater Accra Region of Ghana. *PLoS ONE.* 2013;8(2). <http://journals.plos.org/plosone/article?id=10.1371/journal.pone.0055610>.
15. Poyer S, Shewchuk T, Tougher S, et al. Availability and price of malaria rapid diagnostic tests in the public and private health sectors in 2011: results from 10 nationally representative cross-sectional retail surveys. *Trop Med Int Health.* 2015;20(6):744-756. <https://www.ncbi.nlm.nih.gov/pubmed/25728761>.
16. Prytherch H, Kagoné M, Aninanya GA, et al. Motivation and incentives of rural maternal and neonatal health care providers: a comparison of qualitative findings from Burkina Faso, Ghana and Tanzania. *BMC Health Serv Res.* 2013;13:149. <https://bmchealthservres.biomedcentral.com/articles/10.1186/1472-6963-13-149>.
17. Riley C, Dellicour S, Ouma P, et al. Knowledge and Adherence to the National Guidelines for Malaria Diagnosis in Pregnancy among Health-Care Providers and Drug-Outlet Dispensers in Rural Western Kenya. *Am J Trop Med Hyg.* 2018;98(5):1367-1373. <http://journals.plos.org/plosone/article?id=10.1371/journal.pone.0145616>.
18. The President's Malaria Initiative Ninth Annual Report to Congress. Washington DC: *PMI*; 2015. <http://www.africairs.net/2015/04/pmi-releases-ninth-annual-report/>.
19. WHO: World Malaria Report 2015. Geneva: World Health Organization; 2015. <http://www.who.int/malaria/publications/world-malaria-report-2015/report/en/>.
20. Zurovac D, Githinji S, Memusi D, et al. Major Improvements in the Quality of Malaria Case-Management under the "Test and Treat" Policy in Kenya. *PLOS ONE.* 2014;9(3):e92782. <http://journals.plos.org/plosone/article?id=10.1371/journal.pone.0092782>.