

Think | BIG

Behavior Profile References: Track

1. Githinji S, Kigen S, Memusi D, et al. Using mobile phone text messaging for malaria surveillance in rural Kenya. *Malar J.* 2014;13:107. <https://malariajournal.biomedcentral.com/articles/10.1186/1475-2875-13-107>.
2. Githinji S, Oyando R, Malinga J, et al. Completeness of malaria indicator data reporting via the District Health Information Software 2 in Kenya, 2011–2015. *Malar J.* 2017;16:344. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5561621>.
3. Kabaghe AN, Phiri MD, Phiri KS, van Vugt M. Challenges in implementing uncomplicated malaria treatment in children: a health facility survey in rural Malawi. *Malar J.* 2017;16:419. <https://www.ncbi.nlm.nih.gov/pubmed/29047388>.
4. Karuri J, Waiganjo P, Orwa D, Manya A. DHIS2: The tool to improve health data demand and use in Kenya. *J Health Inform Dev Ctries.* 2014;8(1). <http://www.jhidc.org/index.php/jhidc/article/view/113>. Accessed June 18, 2018.
5. Maina JK, Macharia PM, Ouma PO, Snow RW, Okiro EA. Coverage of routine reporting on malaria parasitological testing in Kenya, 2015–2016. *Glob Health Action.* 2017;10(1):1413266. <https://www.ncbi.nlm.nih.gov/pubmed/29261450>.
6. Mboera LEG, Rumisha SF, Mlacha T, Mayala BK, Bwana VM, Shayo EH. Malaria surveillance and use of evidence in planning and decision making in Kilosa District, Tanzania. *Tanzan J Health Res.* 2017;19(3). <https://www.ajol.info/index.php/thrb/article/view/159505>. Accessed June 18, 2018.
7. Ngwakongni E, Atanga MBS, Quan H. Challenges to implementing a national health information system in Cameroon: Perspectives of stakeholders. *J Public Health Afr.* 2014;5(1). <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5345461>.
8. Okello GA. Producing malaria indicators through district health information software (DHIS2): Practices, processes and challenges in Kenya. [Master's thesis]. *Open University, UK*; 2017. <http://oro.open.ac.uk/52787>.
9. Okello G, Gerrets R, Zakayo S, Molyneux S, Jones C. “Every day they keep adding new tools but they don’t take any away”: Producing indicators for intermittent preventive treatment for malaria in pregnancy (IPTp) from routine data in Kenya. *PLOS ONE.* 2018;13(1):e0189699. <http://journals.plos.org/plosone/article?id=10.1371/journal.pone.0189699>.
10. Phalkey RK, Butsch C, Belesova K, Kroll M, Kraas F. From habits of attrition to modes of inclusion: enhancing the role of private practitioners in routine disease surveillance. *BMC Health Serv Res.* 2017;17:599. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5574140>.
11. Rassi C, Graham K, Mufubenga P, King R, Meier J, Gudoi SS. c. *Malar J.* 2016;15:341. Assessing supply-side barriers to uptake of intermittent preventive treatment for malaria in pregnancy: a qualitative study and document and record review in two regions of Uganda. <https://malariajournal.biomedcentral.com/articles/10.1186/s12936-016-1405-4>.
12. Yukich JO, Butts J, Miles M, et al. A description of malaria sentinel surveillance: a case study in Oromia Regional State, Ethiopia. *Malar J.* 2014;13:88. <https://malariajournal.biomedcentral.com/articles/10.1186/1475-2875-13-88>.