

Think | BIG

Behavior Profile References: Treat

1. ACTwatch Group, Musuva A, Ejersa W, Kiptui R, Memusi D, Abwao E. The malaria testing and treatment landscape in Kenya: results from a nationally representative survey among the public and private sector in 2016. *Malar J.* 2017;16(1). <https://www.ncbi.nlm.nih.gov/pubmed/29268789>.
2. Altaras R, Nuwa A, Agaba B, Streat E, Tibenderana JK, Strachan CE. Why do health workers give anti-malarials to patients with negative rapid test results? A qualitative study at rural health facilities in western Uganda. *Malar J.* 2016;15:23. <https://malariajournal.biomedcentral.com/articles/10.1186/s12936-015-1020-9>.
3. Amoakoh-Coleman M, Agyepong IA, Kayode GA, Grobbee DE, Klipstein-Grobusch K, Ansah EK. Public health facility resource availability and provider adherence to first antenatal guidelines in a low resource setting in Accra, Ghana. *BMC Health Serv Res.* 2016;16:505. <https://bmchealthservres.biomedcentral.com/articles/10.1186/s12913-016-1747-1>.
4. Banchani E, Tenkorang EY. Implementation challenges of maternal health care in Ghana: the case of health care providers in the Tamale Metropolis. *BMC Health Serv Res.* 2014;14:7. <https://bmchealthservres.biomedcentral.com/articles/10.1186/1472-6963-14-7>.
5. Bawate C, Callender-Carter ST, Nsajju B, Bwayo D. Factors affecting adherence to national malaria treatment guidelines in management of malaria among public healthcare workers in Kamuli District, Uganda. *Malar J.* 2016;15:112. <https://malariajournal.biomedcentral.com/articles/10.1186/s12936-016-1153-5>.
6. Boadu NY, Amuasi J, Ansong D, Einsiedel E, Menon D, Yanow SK. Challenges with implementing malaria rapid diagnostic tests at primary care facilities in a Ghanaian district: a qualitative study. *Malar J.* 2016;15:126. <https://malariajournal.biomedcentral.com/articles/10.1186/s12936-016-1174-0>.
7. Burchett HED, Leurent B, Baiden F, et al. Improving prescribing practices with rapid diagnostic tests (RDTs): synthesis of 10 studies to explore reasons for variation in malaria RDT uptake and adherence. *BMJ Open.* 2017;7(3):e012973. <http://bmjopen.bmj.com/content/7/3/e012973>.
8. Febir LG, Baiden FE, Agula J, et al. Implementation of the integrated management of childhood illness with parasitological diagnosis of malaria in rural Ghana: health worker perceptions. *Malar J.* 2015;14:174. <https://malariajournal.biomedcentral.com/articles/10.1186/s12936-015-0699-y>.
9. Hill E, Hess R, Aborigo R, et al. "I don't know anything about their culture": the disconnect between allopathic and traditional maternity care providers in rural northern Ghana. *Afr J Reprod Health.* 2014;18(2):36-45. <https://www.ncbi.nlm.nih.gov/pubmed/25022140>.
10. Johansson EW. Beyond 'test and treat' – malaria diagnosis for improved pediatric fever management in sub-Saharan Africa. *Glob Health Action.* 2016;9(1):31744. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5165054>.
11. Kabaghe AN, Phiri MD, Phiri KS, van Vugt M. Challenges in implementing uncomplicated malaria treatment in children: a health facility survey in rural Malawi. *Malar J.* 2017;16:419. <https://www.ncbi.nlm.nih.gov/pubmed/29047388>.
12. Kwarteng A, Asante KP, Abokyi L, et al. Provider compliance to artemisinin-based combination therapy at primary health care facilities in the middle belt of Ghana. *Malar J.* 2015;14:361. <https://malariajournal.biomedcentral.com/articles/10.1186/s12936-015-0902-1>.

13. Mannan AA, Elmardi KA, Idris YA, Spector JM, Ali NA, Malik EM. Do frontline health care providers know enough about artemisinin–based combination therapy to rationally treat malaria? A cross-sectional survey in Gezira State, Sudan. *Malar J.* 2015;14:131. <https://malariajournal.biomedcentral.com/articles/10.1186/s12936-015-0652-0>.
14. Mbonye AK, Buregyeya E, Rutebemberwa E, et al. Treatment and prevention of malaria in pregnancy in the private health sector in Uganda: implications for patient safety. *Malar J.* 2016;15:212. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4831190>.
15. Michael D, Mkunde SP. The malaria testing and treatment landscape in mainland Tanzania, 2016. *Malar J.* 2017;16:202. <https://malariajournal.biomedcentral.com/articles/10.1186/s12936-017-1819-7>.
16. Nwameme AU, Phillips JF, Adongo PB. Compliance with Emergency Obstetric Care Referrals Among Pregnant Women in an Urban Informal Settlement of Accra, Ghana. *Matern Child Health J.* 2014;18(6):1403-1412. <https://www.ncbi.nlm.nih.gov/pubmed/24178157>.
17. Oduro-Mensah E, Kwamie A, Antwi E, et al. Care Decision Making of Frontline Providers of Maternal and Newborn Health Services in the Greater Accra Region of Ghana. *PLoS ONE.* 2013;8(2). <http://journals.plos.org/plosone/article?id=10.1371/journal.pone.0055610>.
18. Phanalasy S. The malaria testing and treatment landscape in the southern Lao People's Democratic Republic (PDR). *Malar J.* 2017;16:169. <https://malariajournal.biomedcentral.com/articles/10.1186/s12936-017-1769-0>.
19. Prytherch H, Kagoné M, Aninanya GA, et al. Motivation and incentives of rural maternal and neonatal health care providers: a comparison of qualitative findings from Burkina Faso, Ghana and Tanzania. *BMC Health Serv Res.* 2013;13:149. <https://bmchealthservres.biomedcentral.com/articles/10.1186/1472-6963-13-149>.
20. Riley C, Dellicour S, Ouma P, et al. Knowledge and Adherence to the National Guidelines for Malaria Diagnosis in Pregnancy among Health-Care Providers and Drug-Outlet Dispensers in Rural Western Kenya. *Am J Trop Med Hyg.* 2018;98(5):1367-1373. <http://journals.plos.org/plosone/article?id=10.1371/journal.pone.0145616>.
21. Salomão CA, Sacarlal J, Chilundo B, Gudo ES. Prescription practices for malaria in Mozambique: poor adherence to the national protocols for malaria treatment in 22 public health facilities. *Malar J.* 2015;14:483. <https://www.ncbi.nlm.nih.gov/pubmed/26628068>.
22. The President's Malaria Initiative Ninth Annual Report to Congress. Washington DC: *PMI*; 2015. [https://www.pmi.gov/docs/default-source/default-document-library/tools-curricula/pmi-technical-guidance-\(march-2016\).pdf](https://www.pmi.gov/docs/default-source/default-document-library/tools-curricula/pmi-technical-guidance-(march-2016).pdf).
23. Umeano-Enemuoh JC, Uzochukwu B, Ezumah N, Mangham-Jefferies L, Wiseman V, Onwujekwe O. A qualitative study on health workers' and community members' perceived sources, role of information and communication on malaria treatment, prevention and control in southeast Nigeria. *BMC Infect Dis.* 2015;15:437. <https://www.ncbi.nlm.nih.gov/pubmed/26493559>.
24. WHO: World Malaria Report 2017. Geneva: World Health Organization; 2017. <http://www.who.int/malaria/media/world-malaria-report-2017/en/#Diagnostic%20testing%20and%20treatment>.
25. Yoder PS, Nsabagasani X, Eckert E, Moran A, Yé Y. Perspectives of health care providers on the provision of intermittent preventive treatment in pregnancy in health facilities in Malawi. *BMC Health Serv Res.* 2015;15:354. <https://bmchealthservres.biomedcentral.com/articles/10.1186/s12913-015-0986-x>.