

Think | **BIG**

Behavior Profile References: Comprehensive Post-Natal Care

1. Adam T, Lim SS, Mehta S, et al. Achieving the millennium development goals for health cost effectiveness analysis of strategies for maternal and neonatal health in developing countries. *BMJ: British Medical Journal*. 2005(7525):1107.
2. Ahmed S, Sobhan F, Islam A, Barkat-e-Khuda n. Neonatal morbidity and care-seeking behaviour in rural Bangladesh. *J Trop Pediatr*. 2001;47(2):98-105.
3. Bassat Q, Simkiss D, Edmond K, Bose A, Troy S, eds. Care of the newborn in developing countries book 2 chapter 1 problems of the newborn. *Journal of Tropical Pediatrics*.
4. Callaghan-Koru JA, Seifu A, Tholandi M, et al. Newborn care practices at home and in health facilities in 4 regions of Ethiopia. *BMC pediatrics*. 2013;13(1):198.
5. Carlo WA, Goudar SS, Jehan I, et al. Newborn-care training and perinatal mortality in developing countries. *New England Journal of Medicine*. 2010;362(7):614-623.
6. Darmstadt GL, Hassan M, Balsara ZP, Winch PJ, Gipson R, Santosham M. Impact of clean delivery-kit use on newborn umbilical cord and maternal puerperal infections in Egypt. *J Health Popul Nutr*. 2009;27(6):746-754.
7. Fikree FF, Ali TS, Durocher JM, Rahbar MH. Newborn care practices in low socioeconomic settlements of Karachi, Pakistan. *Soc Sci Med*. 2005;60(5):911-921.
8. Fujino Y, Sasaki S, Igarashi K, et al. Improvement in mothers' immediate care-seeking behaviors for children's danger signs through a community-based intervention in Lusaka, Zambia. *The Tohoku Journal of Experimental Medicine*. 2009;217(1):73-85.
9. Ganle JK, Parker M, Fitzpatrick R, Otupiri E. A qualitative study of health system barriers to accessibility and utilization of maternal and newborn healthcare services in Ghana after user-fee abolition. *BMC pregnancy and childbirth*. 2014;14(1):425.
10. Kayode, Gbenga A., et al. Individual and community determinants of neonatal mortality in Ghana: a multilevel analysis. *BMC pregnancy and childbirth*. 14.1 (2014): 165.
11. Marsh DR, Darmstadt GL, Moore J, Daly P, Oot D, Tinker A. Advancing newborn health and survival in developing countries: A conceptual framework. *Journal of perinatology*; Document Updated: June 2018. official journal of the California Perinatal Association. 2002;22(7):572-576.
12. MEASURE Evaluation. Newborn health:
http://www.cpc.unc.edu/measure/prh/rh_indicators/specific/nb
13. Murray J, Adeyi GN, Graeff J, et al. Emphasis behaviors in maternal and child health: Focusing on caretaker behaviors to develop maternal and child health programs in communities. 1997.
14. Newbrander W, Natiq K, Shahim S, Hamid N, Skena NB. Barriers to appropriate care for mothers and infants during the perinatal period in rural Afghanistan: A qualitative assessment. *Global public health*. 2014;9 Suppl 1(sup1):S109.
15. Shiferaw S, Spigt M, Godefrooij M, Melkamu Y, Tekie M. Why do women prefer home births in Ethiopia? *BMC pregnancy and childbirth*. 2013;13(1):5.

16. Sinha LN, Kaur P, Gupta R, Dalpath S, Goyal V, Murhekar M. Newborn care practices and home-based postnatal newborn care programme - Mewat, Haryana, India, 2013. *Western Pacific surveillance and response journal: WPSAR*. 2014;5(3):22-29.
17. Sreeramareddy CT, Joshi HS, Sreekumaran BV, Giri S, Chuni N. Home delivery and newborn care practices among urban women in western Nepal: A questionnaire survey. *BMC pregnancy and childbirth*. 2006;6(1):27.
18. World Health Organization. WHO recommendations on postnatal care of the mother and newborn. *World Health Organization*. 2014.
19. WHO, UNICEF. Home visits for the newborn child: A strategy to improve survival. 2009.
20. WHO. Newborn death and illness. WHO Web site:
http://www.who.int/pmnch/media/press_materials/fs/fs_newbornddeath_illness/en/
21. WHO. Newborns: Reducing mortality. WHO Web site.
<http://www.who.int/mediacentre/factsheets/fs333/en>