

Think | **BIG**

Behavior Profile References: Insecticide-Treated Net Use

1. Ahmed SM, Zerihun A. Possession and usage of insecticidal bed nets among the people of Uganda: Is BRAC Uganda health programme pursuing a pro-poor path? *PLoS One*. 2010;5(9):e12660. Accessed Sep 26, 2016. doi: 10.1371/journal.pone.0012660.
2. Awosan K, Ibrahim MTO, Alayandae MO, Isah BA, Yunusa E, Mahmud MB. Prevalence and barriers to the use of insecticide treated nets among pregnant women attending ante-natal clinic at specialist hospital Sokoto, Nigeria. *Journal of Public Health and Epidemiology*. 2013.
3. Bennett A, Smith SJ, Yambasu S, et al. Household possession and use of insecticide-treated mosquito nets in Sierra Leone 6 months after a national mass-distribution campaign. *PLoS One*. 2012;7(5):e37927. Accessed Sep 26, 2016. doi: 10.1371/journal.pone.0037927.
4. Comfort AB, Krezanoski PJ. The effect of price on demand for and use of bednets: evidence from a randomized experiment in Madagascar. *Health Policy Plan*. 2017;32(2):178-193. doi:10.1093/heapol/czw108.
5. Communications Support for Health, (CSH). Zambia communications support for health: Stop malaria champion communities programme evaluation. 2014.
6. Danhoundo G, Wiktorowicz ME, Premji SS, Nasiri K. Determinants of bed net policy implementation: A case study of Southern Benin. *The International Journal of Health Planning and Management*. 33(1):e279-e292. doi:10.1002/hpm.2465
7. Gemade EI, Earland J. Long lasting insecticidal nets (LLINs) ownership and use: A qualitative study to explore why people in Kuje area council of federal capital territory of Nigeria are not sleeping under the LLINs for malaria prevention. *The Journal of MacroTrends in Applied Science*. 2013;1(1):26-41.
8. Gikandi PW, Noor AM, Gitonga CW, Ajanga AA, Snow RW. Access and barriers to measures targeted to prevent malaria in pregnancy in rural Kenya. *Tropical Medicine & International Health*. 2008;13(2):208-217. doi: 10.1111/j.1365-3156.2007.01992.x.
9. Godpower MC, Ibrahim A, Grema BA. Knowledge of malaria and adherence to its preventive measures among adults attending out-patient clinics of a Nigerian tertiary hospital: Has anything changed? *African Journal of Medical and Health Sciences*. 2017;16(1):43-51.
10. Kanyangarara M, Hamapumbu H, Mamini E, et al. Malaria knowledge and bed net use in three transmission settings in southern Africa. *Malaria Journal*. 2018;17:41. doi:10.1186/s12936-018-2178-8
11. Katz I, Komatsu R, Low-Beer D, Atun R. Scaling up towards international targets for AIDS, tuberculosis, and malaria: Contribution of global fund-supported programs in 2011–2015. *PLoS One*. 2011;6(2):e17166. Accessed Sep 26, 2016. doi: 10.1371/journal.pone.0017166.
12. Koenker H, Yukich JO. Effect of user preferences on ITN use: a review of literature and data. *Malaria Journal*. 2017;16:233. doi:10.1186/s12936-017-1879-8.
13. Kolaczinski JH, Kolaczinski K, Kyabayinze D, et al. Costs and effects of two public sector delivery channels for long-lasting insecticidal nets in Uganda. *Malar J*. 2010;9:102. Accessed Sep 26, 2016. doi: 10.1186/1475-2875-9-102.

14. Kweku M, Webster J, Taylor I, Burns S, Dedzo M. Public-private delivery of insecticide-treated nets: A voucher scheme in Volta region, Ghana. *Malaria Journal*. 2007;6(1):14.
15. Lam Y, Harvey SA, Monroe A, et al. Decision-making on intra-household allocation of bed nets in Uganda: Do households prioritize the most vulnerable members? *Malaria Journal*. 2014;13(1):183. doi: 10.1186/1475-2875-13-183.
16. Larsen DA, Keating J, Miller J, et al. Barriers to insecticide-treated mosquito net possession 2 years after a mass free distribution campaign in Luangwa district, Zambia. *PLoS One*. 2010;5(11):e13129. Accessed August 30, 2016. doi: 10.1371/journal.pone.0013129.
17. Makungu C, Stephen S, Kumburu S, et al. Informing new or improved vector control tools for reducing the malaria burden in Tanzania: a qualitative exploration of perceptions of mosquitoes and methods for their control among the residents of Dar es Salaam. *Malaria Journal*. 2017;16:410. doi:10.1186/s12936-017-2056-9.
18. Malaria Communities Program. Achieving, tracking, and maintaining high ITN coverage: Community strategies. 2013.
19. Marchant T, Schellenberg D, Nathan R, et al. Assessment of a national voucher scheme to deliver insecticide-treated mosquito nets to pregnant women. *CMAJ*. 2010;182(2):152-156. Accessed Sep 26, 2016. doi: 10.1503/cmaj.090268.
20. Monroe A, Harvey SA, Lam Y, et al. "People will say that I am proud": A qualitative study of barriers to bed net use away from home in four Ugandan districts. *Malaria journal*. 2014;13(1):82. doi: 10.1186/1475-2875-13-82.
21. Msellemu D, Shemdoe A, Makungu C, et al. The underlying reasons for very high levels of bed net use, and higher malaria infection prevalence among bed net users than non-users in the Tanzanian city of Dar es Salaam: a qualitative study. *Malaria Journal*. 2017;16:423. doi:10.1186/s12936-017-2067-6
22. Njoroge FK, Kimani VN, Ongore D, Akwale WS. Use of insecticide treated bed nets among pregnant women in Kilifi district, Kenya. *East African Medical Journal*. ;86(7):314-22. Accessed August 30, 2016. doi: 10.4314/eamj.v86i7.54145.
23. Oladimeji KE, Tsoka. Gwegweni JM, Gengiah S, Daftary A, Naidoo K. Barriers to effective uptake of malaria prevention interventions in Ibadan, South West Nigeria: a qualitative study. *International Journal Of Community Medicine And Public Health*. 2018;5(4):1304-1310.
24. Pell C, Straus L, Andrew EVW, Meñaca A, Pool R. Social and cultural factors affecting uptake of interventions for malaria in pregnancy in Africa: A systematic review of the qualitative research. *PLoS One*. 2011;6(7):e22452. Accessed Sep 26, 2016. doi:10.1371/journal.pone.0022452.
25. Pettifor A, Taylor E, Nku D, et al. Free distribution of insecticide treated bed nets to pregnant women in Kinshasa: An effective way to achieve 80% use by women and their newborns. *Trop Med Int Health*. 2009;14(1):20-28. Accessed Sep 26, 2016. doi: 10.1111/j.1365-3156.2008.02179.x.
26. Rhee M, Sissoko M, Perry S, McFarland W, Parsonnet J, Doumbo O. Use of insecticide-treated nets (ITNs) following a malaria education intervention in Piron, Mali: A control trial with systematic allocation of households. *Malaria journal*. 2005;4(1):35.
27. Rickard DG, Dudovitz RN, Wong MD, et al. Closing the gap between insecticide treated net ownership and use for the prevention of malaria. *Prog Community Health Partnersh*. 2011;5(2):123-131. Accessed Sep 26, 2016. doi: 10.1353/cpr.2011.0018.

28. Salomão C, Sacarlal J, Gudo ES. Assessment of coverage of preventive treatment and insecticide-treated mosquito nets in pregnant women attending antenatal care services in 11 districts in Mozambique in 2011: the critical role of supply chain. *Malaria Journal*. 2017;16:223. doi:10.1186/s12936-017-1872-2
29. Singh M, Brown G, Rogerson SJ. Ownership and use of insecticide-treated nets during pregnancy in sub-Saharan Africa: A review. *Malar J*. 2013;12:268. Accessed August 30, 2016. doi: 10.1186/1475-2875-12-268.
30. Stewart T, Marchand RP. Factors that affect the success and failure of insecticide treated net programs for malaria control in SE assai and the western pacific. WHO-WPRO.
31. Tesfa G. Peoples' belief, attitude, and practice in the use of insecticide treated bed net (ITN): The case of Serbo, Nada, and Asendabo towns, Jimma zone, southwest Ethiopia. *Ethiopian Journal of Education and Sciences*. 2012;8(1):93-106. Accessed August 30, 2016.
32. Thompson LH. Head of household characteristics influencing insecticide treated net use in Ghana. *ProQuest Dissertations Publishing*; 2014.
33. West PA, Protopopoff N, Rowland MW, et al. Evaluation of a national universal coverage campaign of long-lasting insecticidal nets in a rural district in north-west Tanzania. *Malaria Journal*. 2012;11(1):273. doi: 10.1186/1475-2875-11-273.
34. WHO. World malaria report 2011. WHO. 2011. Accessed Sep 26, 2016.